

OPEN ACCESS –RESEARCH ARTICLE

Local dominance of exotic plants declines with residence time: a role for plant soil feedback?

Tanja A.A. Speek^{1,2,3,*}, Joop H.J. Schaminée^{4,5}, Jeltje M Stam⁶, Lambertus A.P. Lotz¹, Wim A. Ozinga⁴ and Wim H. van der Putten^{2,3}

¹ Wageningen University & Research Centre, Plant Research International, Droevendaalsesteeg 1, 6708 PB Wageningen, Netherlands,

² Wageningen University & Research Centre, Laboratorium of Nematology, Droevendaalsesteeg 1, 6708 PB Wageningen, Netherlands,

³ Netherlands Institute of Ecology (NIOO-KNAW), Department of Terrestrial Ecology, Droevendaalsesteeg 10, 6708 PB Wageningen, Netherlands,

⁴ Wageningen University & Research Centre, Centre Ecosystem Studies, Droevendaalsesteeg 3a, 6708 PB Wageningen, Netherlands,

⁵ Radboud University Nijmegen, Department Ecology, Aquatic Ecology & Environmental Biology Research Group, Heyendaalseweg 135, 6525 AJ Nijmegen, Netherlands,

⁶ Wageningen University & Research Centre, Entomology, Droevendaalsesteeg 1, 6708 PB Wageningen, Netherlands,

*Correspondence: E-mail: tanjaspeek@gmail.com

Received: 3 June 2014; Revised: 25 August 2014; Accepted: 20 January 2015

Published by Oxford University Press on behalf of the Annals of Botany Company. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

ABSTRACT

Recent studies have shown that introduced exotic plant species may escape from their native soil-borne pathogens, but that they become exposed to increased soil pathogen activity in the new range when time since introduction increases. Other studies have shown that introduced exotic plant species become less dominant when time since introduction increases, and that plant abundance may be controlled by soil-borne pathogens, however, no study yet has tested if these soil effects might explain the decline in dominance of exotic plant species following their initial invasiveness. Here we determine plant-soil feedback of 20 plant species that have been introduced into The Netherlands. We tested the hypotheses that (1) exotic plant species with a longer residence time have a more negative soil feedback, and (2) greater local dominance of the introduced exotic plant species correlates with less negative, or more positive plant-soil feedback. Although the local dominance of exotic plant species decreased with time since introduction, there was no relationship of local dominance with plant-soil feedback. Plant-soil feedback also did not become more negative with increasing time since introduction. We discuss why our results may deviate from some earlier published studies and why plant-soil feedback may not in all cases, or not in all comparisons explain patterns of local dominance of introduced exotic plant species.

Keywords: Exotic species; introduced species; local dominance; macro ecology; residence time; soil-borne enemy.

INTRODUCTION

An important challenge for invasion ecologists is to predict the course of invasions of introduced exotic species. This requires insight in the factors that may control the abundance and dominance of species in both their native and new ranges. It has been well established that regional distribution of exotic plant species increases with residence time (Pyšek *et al.* 2004; Hamilton *et al.* 2005; Wilson *et al.* 2007; Milbau and Stout 2008; Bucharova and van Kleunen 2009; Gassó *et al.* 2009). It has also been argued that increased residence time may result in lower local dominance and invasiveness (Carpenter and Cappuccino 2005; Hawkes 2007; Speek *et al.* 2011). Local dominance of introduced exotic plant species may be, at least in part, driven by interactions with soil biota, including effects of soil-borne enemies and symbionts (Inderjit and Van der Putten 2011). The question that we address in the present study is how residence time and local dominance of exotic plant species may relate to enemy impact of the soil biota. Ultimately, this information may be used to enhance predictions on the course of invasiveness of introduced exotic plant species.

A possible explanation for lower local dominance of introduced exotic plant species with a long residence time is that enemy species may increasingly adapt and accumulate when time of exposure to the new hosts increases (Diez *et al.* 2010; Hawkes 2007; Dostal *et al.* 2013). Both aboveground (Bentley and Whittaker 1979; Gange and Brown 1989) and belowground (Klironomos 2002; Johnson *et al.* 2012; Mangan *et al.* 2010; van der Putten *et al.* 1993) enemies may control local plant dominance. Release from natural enemies by introduction to a new range has been proposed to

enhance the performance of species and, therefore, their invasiveness (Elton 1958; Keane and Crawley 2002). This 'enemy release hypothesis' (Keane and Crawley 2002) has been supported by surveys showing that introduced plant species have fewer enemies in their novel than native range (e.g. Mitchell and Power 2003).

Thus far, the majority of research on enemy release of exotic plant species has been dedicated to aboveground enemies. However, an increasing amount of studies is showing that introduced exotic plant species can be released from native soil-borne enemies as well (Reinhart *et al.* 2003; Reinhart *et al.* 2010; Callaway *et al.* 2004; Gundale 2014). Introduced exotic plant species suffer less from soil-enemies of the invaded range than congeners that are native in that range (Maron and Vilà 2001; Agrawal *et al.* 2005; Engelkes *et al.* 2008).

The change in performance of exotic species with progressing residence time has been described for several invaders (Simberloff and Gibbons 2004). Loss of exotic dominance might be caused by evolutionary adaptation of enemies in the new range to the introduced plant species (Müller-Schärer *et al.* 2004). Such adaptive potential may be deduced from reported higher frequencies of specialist compared to generalist herbivores (Andow and Imura 1994), higher exposure (Mitchell *et al.* 2010), and higher impact (Hawkes 2007) of enemies on crop and exotic plant species in relation to increasing residence time. Similarly, in New Zealand plant-soil feedback of 12 exotic plant species related negatively to their residence time (Diez *et al.* 2010) and in the Czech Republic giant hogweed (*Heracleum mantegazzianum*) developed negative feedback effects from the soil biota in

fields that had been colonized for some decades (Dostál *et al.* 2013). However, in these latter studies, increased enemy exposure has not yet been related to local dominance of the exotic plant species, which is the key aim of the present study.

A recent analysis established that exotic plant species with a long residence time in the Netherlands have lower local dominance than recently introduced species (Speek *et al.* 2011). Until now, no study has related such patterns in local dominance to plant-soil feedback effects. Therefore, in the present study, we determine how residence time, local dominance and soil pathogen effects to exotic species may relate to each other. We tested soil pathogen effects by plant-soil feedback approach (Bever *et al.* 1997), which is a way to experimentally integrate all positive and negative interactions between plants and the soil biota. We first tested the hypothesis that species with a longer residence time have a more negative plant-soil feedback (Diez *et al.* 2010). Then, we tested the hypothesis that species with a more positive plant soil feedback have a higher local dominance (Klironomos 2002).

METHODS

Data on plants, their residence time and local dominance

Data on residence time were derived from information on period of naturalization according to the standard list of the Dutch flora (Tamis *et al.* 2004). Data on local dominance were derived from the Dutch Vegetation Database (Schaminée *et al.* 2007), containing over 500,000 vegetation records including data on local species cover in plots varying from 1 by 1 m² to 10 by 10 m². Plot sizes used for recording depended on the characteristics

of vegetation, for example largest plot sizes were used for forests. Data on plant species cover were used to calculate local dominance as [the number of vegetation records with that species having > 10 per cent ground cover / the total number of vegetation records with that plant species] x 100 % (Speek *et al.* 2011). Therefore, local dominance expresses the frequency of how often a plant species has a minimum cover of ten percent, when present in the vegetation record. In order to exclude recorder bias, for example due to avoiding taking records of vegetation heavily invaded by exotic plant species, we used expert judgment to check and where necessary adjust the cover data (Speek *et al.* 2011).

Soil feedback experiment

We used a selection of 20 introduced exotic plant species in the Netherlands for a plant-soil feedback experiment (Supplements). The selection of 20 plant species was based on a number of criteria. First, we excluded woody species, because the length of the plant-soil feedback is too limited for capturing a substantial part of the life cycle of trees. We then selected as many as possible plant species from riverine areas in order to be able to use the same soil origin for all plant species. Finally, the selection was limited as the seeds of some plant species did not germinate. Seeds had been collected by specialized seed companies that collect seeds locally, or by ourselves or colleagues.

Of the 20 plants species, 14 occur in the Millingerwaard (Dirkse *et al.* 2007), a riverine floodplain area of 700 hectares. Millingerwaard is a nature reserve in the riverine floodplain of the river Waal, which is the southern branch of the Rhine river in the Netherlands (51°87' N, 6°01' E). Three other

species occur near or in other riverine areas in the Netherlands and the remaining three occur outside riverine areas. We collected soil from the Millingerwaard area, instead of from a larger variety of sites, as soil from a variety of sites would have introduced additional variation due to soil type, fertility, pH etc. All plant species were forbs that varied in local dominance from 5 - 38 per cent and in residence time from 75 to 400 years.

Seeds were germinated on glass beads placed in demineralised water. Germination was carried out in transparent plastic containers of 17 x 12 x 5 cm that were placed under conditions of 16 hrs 22 °C in the light (day) and 8 hrs 10 °C in the dark (night). *Xanthium strumarium* seeds were germinated at a higher temperature: 16 hrs 32 °C and 8 hrs 20 °C. Germinated seedlings were stored at 4 °C and 10/14 hrs light/dark until transplantation in soil, to ensure equal sizes at start of the experiment. Soil was collected from five random locations in Millingerwaard. Soil to be used as inoculum was collected in October 2010, prior to the first phase of the experiment. Soil from the five sampling locations was sieved (mesh size 5 mm) to remove coarse roots, stones and other large particles, and subsequently homogenized. The bulk soil was collected in January 2010, sterilized by gamma irradiation (25 KGray) and stored in sealed plastic bags at 4 °C until use.

The sensitivity of exotic plant species to soil-borne enemies was determined in a so-called two phase plant-soil feedback experiment (Bever *et al.* 1997). In the first phase, which started from one pooled sample, the seedlings were grown to condition the field soil. In that phase, soil biota that can grow on resources provided by that particular plant species are enumerated (Grayston *et al.* 1998; Kowalchuk *et al.* 2002). In the second

phase, we kept all replicates of own soil separate. In order to do so, the soil of each pot was split in two halves: one half was used as own soil, whereas the other half was mixed with the halves of all other replications and species, to be used as away soils. The replicates of the mixed soil were not kept intact, because there was no relationship between replicate 1 conditioned by species A or B. Comparing plant performances in own and mixed soils enabled us to make a home (own) versus away (mixed) comparison, which is a less sensitive and ecologically more realistic method of detecting plant-soil feedback effects than a comparison of non-sterilized versus sterilized soil (Kulmatiski *et al.* 2008). In the final analysis, plant species was the unit of replication.

For the first –conditioning– phase, bulk soil and inoculum were mixed at a ratio of 4:1, with a total of 1200 gram soil per pot on a dry weight basis. Pots of 1.3 L were used. For the second –feedback– phase, ‘own soil’ and ‘mixed soil’ were homogenized with sterilised bulk soil at a ratio of 1:1 in order to keep pot volumes equal between the two feedback phases. For each plant species, we had five independent replicates with own and five with mixed soil. Every pot contained three seedlings, except *Amaranthus retroflexus* that was planted as two seedlings per pot due to poor germination of the seeds. Dead seedlings were replaced until the first week after transplanting. Greenhouse conditions were maintained at 60% RH, day temperature 21 °C, night temperature 16 °C. Daylight was supplemented with lamps (SON-T Agro, 225 $\mu\text{mol}^{-1} \text{m}^{-2}$), to ensure a minimum of 16 hours light per day.

Before planting, the water content in each pot was set at 20% (w/w). Plants were supplied with water three times a week and once a week the

water content was re-set to 20% by weighing. Plants received 10 ml 0.5 strength Hoagland per pot in weeks two, three and four, and 20 ml in weeks five and six after transplanting in order to meet increasing demand. Plants were harvested six weeks after planting. The length of growth was the same for both phases, which is relatively short, but ample for testing feedback responses (Van der Putten *et al.* 1988). When harvesting, shoots of the three (or two) plants per pot were clipped at ground level, pooled, dried in paper bags at 75 °C until constant weight and weighed, so that biomass data per pot were obtained.

Statistical analysis

The effect of soil feedback on shoot and root biomass was calculated as $\ln[(\text{biomass in own soil})/(\text{biomass in mix soil})]$ (Brinkman *et al.* 2010). We assigned pairs of own soil with mixed soil randomly. To analyze if residence time or local dominance could explain mean shoot and root feedback responses we used linear models. The unit of replication was the plant species. For residence time we used models with a normal distribution, for local dominance we used models with a binomial distribution and a logit link, with binomial totals set to 50% (the highest value in our dataset).

We analyzed which traits and other factors related best to residence time by a model selection procedure within a linear model with a normal distribution. Thus, we selected the best minimal adequate model with the lowest Akaike Information Criterion value from all possible subsets. Although time and dominance were related, the relation of a trait or other factor to residence time may not necessarily imply that there is a relation with local dominance as well. Therefore, the factors in the best minimal adequate model

were added to a generalized linear model with residence time explaining local dominance. By adding each factor separately, we analyzed which one significantly changed the model. Factors that affected the model were likely to be a better explanation for variation in local dominance than residence time. For explaining local dominance we used a binomial distribution with a logit link, binomial totals set at 50 and accounting for overdispersion. All analysis were done in Genstat version 14.

RESULTS

Opposite to our hypothesis, we found neither a significant relationship between residence time and plant-soil feedback of the exotic plant species, nor for shoots ($F=0.10$, $t_{18}=-0.32$, $p=0.751$, Fig. 1) nor roots ($F=0.41$, $t_{18}=-0.64$, $p=0.529$). Local plant dominance also did not relate to the feedback effect on shoots ($F=0.09$, $t_{18}=-0.31$, $p=0.763$) or roots ($F=0.73$, $t_{18}=-0.85$, $p=0.404$).

Excluding species from riverine habitats, which may not be responsive to soil biota from that habitat, or Fabaceae species, which may have a different feedback due to symbiosis with rhizobia did not alter the significance of the results (data not shown). Therefore, our hypotheses that species with a longer residence time have a more negative plant-soil feedback, and that species with a less negative or more positive plant-soil feedback have a higher local dominance were not supported.

DISCUSSION

In our study we tested the hypotheses that species with a longer residence time have a more negative plant-soil feedback and that species with a less

negative, or more positive plant soil feedback have a higher local dominance. We used an experimental approach to measure soil-borne enemy impact by plant-soil feedback approach. However, opposite to a study from New Zealand (Diez *et al.* 2010), and to a study on introduced *H. mantegazzianum* in the Czech Republic (Dostál *et al.* 2013) we did not find such a relationship between time since introduction of 20 exotic plant species in the Dutch flora and plant-soil feedback.

There are several possible explanations for these results. Our results could indicate that not all introduced exotic plant species develop negative plant-soil feedback when time since introduction increases. In the field, other ecological processes may be influencing community composition and aboveground interactions can either increase or decrease with the strength of the belowground interactions. Another possible explanation concerns the choice of soils for the plant-soil feedback experiment. We have chosen soils from areas where most exotic plant species may occur, but we did not use soils from the root zone of particular populations. This approach has led to marked differences in plant-soil feedback between natives and exotics (Van Grunsven *et al.* 2007, Engelkes *et al.* 2008), however, it has resulted in scattered results when testing soil responses across an entire native range (Van Grunsven *et al.* 2010).

The results may also depend on the relatively short conditioning and testing phases of 6 weeks each. Test phases of 6 weeks can detect feedback effects (Van der Putten *et al.* 1988). Longer test periods may even result in pot limitations, which may obfuscate results. Conditioning for 6 weeks will

have been relatively short, but to our experience this is possible when adding soil inocula to sterilized soil, as has been done in the present study.

Our use of pooled soils as 'away' treatment may have provided a conservative estimate of plant-soil feedback effects, because of reducing variances. Nevertheless, since we did not find significant relationships with time since abandonment, or local dominance, our results show that even with a highly sensitive test still no relationship could be detected between time since introduction, or local dominance, and plant-soil feedback. Mixing soils from all plant species to produce 'away' soils could theoretically have led to single pathogens dominating the entire away soil community. However, a previous addition study using a variety of amounts of soil inocula showed that soil feedback effects increased gradually with the amount of inoculum added (Van der Putten *et al.* 1988), which does not point at a disproportional role of pathogens from single plant species in the away soil mixtures.

Plant-soil biota interactions are highly local (Levine *et al.* 2006; Bezemer *et al.* 2010; Genung *et al.* 2012), and adaptation of soil organisms to new plant species does not take place at a national, but at a local scale through direct interactions between plant roots and the soil biota (Schweitzer *et al.* 2008; Lankau *et al.* 2009; Lau and Lennon 2012; Lau and Lennon 2011). As the feedback was estimated at a regional scale, also the local dominance was measured at a regional scale (first occurrence in the Netherlands). Using first occurrence in a larger region as estimate of residence time could result in an over-estimation of the local residence time. On the other hand, the study from New Zealand (Diez *et al.* 2010) also used

data on residence time for the entire country and not specifically for the sites from which the soil has been collected.

We expected plant-soil feedback to be negatively related to local dominance (Klironomos 2002; Mangan *et al.* 2010). However, in our study we did not observe such an inverse relationship. A possible explanation is that the previous studies by Klironomos (2002) and Mangan *et al.* (2010) on dominance-feedback relationships have been based on native species, and that these relations may differ when considering exotic species. Moreover, we used dominance estimates averaged across the entire Netherlands (Speek *et al.* 2011), which differs from the local dominance estimates as used in other studies (e.g. Klironomos 2002). National estimates (in the case of the Netherlands concerning an area of appr. 150 x 300 km) will not provide accurate information about the local dominance of exotics in the riverine ecosystem where the soil for testing plant-soil feedback originated from. Therefore, it is possible that soil origin and plant dominance data were not well linked to each other, or that a relationship between plant-soil feedback and dominance works out differently for exotic plant species than for natives. Alternatively, our study may add to other examples where plant dominance does not relate to plant-soil feedback (Reinhart 2012).

An alternative explanation for the rejection of our hypotheses could be that the evolutionary dynamics leading to increased enemy pressure on exotic plant species is not strong enough to result in a change in mean local dominance. Meta-analyses have shown that a general pattern of decreased enemy numbers on exotic species in the novel range was not reflected in a general pattern of higher plant performance (Chun *et al.* 2010). Adaptation

can occur both at the soil species level but also at the plant species level. This adaptation at two fronts is likely to result in a mixed general outcome. Moreover, while local dominance has been assumed to increase after introduction to a new range (Keane and Crawley 2002), recent work has shown that most species have the same dominance in both their introduced and native ranges (Firn *et al.* 2011). Clearly, local dominance is a complex trait, with a high variation both between and within species that can be influenced by a large number of ecological processes.

CONCLUSIONS

We found no support for the hypothesis that the negative relationship between residence time and local dominance of exotic species in The Netherlands is caused by an increase in negative plant soil feedback. It may be that data on residence time, dominance, enemy exposure and impact need to be collected all from the same area, or that different choices in plant-soil feedback approach need to be made (e.g. longer conditioning and/or feedback phases, a more sensitive 'away' soil treatment). Alternatively, it might be better to track single species across an introduction gradient (Lankau *et al.* 2009; Lankau 2011). It could also mean that not all introduced exotic plant species develop negative plant-soil feedback when time since introduction increases or that the hypothesized effect of increasing enemy pressure on dominance of introduced exotic plant species might not be strong enough to emerge from examining a large diversity of species across a variation of locations. Therefore even though we are aware of weaknesses of our paper (aspects of the experimental design that were not ideal, for example sampling of soil from

one location that did not include all of the study species, pooling "away" soils, method of pairing of home and away pots to calculate response ratios), our results may add to the debate on change in invasiveness of exotic plant species after introduction.

SUPPORTING INFORMATION

The following table is available in the online version of this article:

File 1. Table. Lists the species that were included in the experiment and data on their local dominance and residence time in the Netherlands.

SOURCES OF FUNDING

TAA Speek and LAP Lotz were funded by the former Dutch Ministry of Agriculture, Nature and Food Quality, FES-programme 'Versterking Infrastructuur Plantgezondheid'. W van der Putten was supported by ALW-Vici grant. WA Ozinga was supported by the Dutch Science Foundation (NWO Biodiversity Works).

CONTRIBUTIONS BY AUTHORS

All authors contributed to the experimental set up, TAA Speek, JM Stam and WvdPutten performed the experiment, TAA Speek performed all analyses, JHJ Schaminee and WA Ozinga provided data on local dominance of plant species, TAA Speek and WvdPutten wrote the paper.

CONFLICTS OF INTEREST

No conflicts of interest

ACKNOWLEDGEMENTS

We thank Staatsbosbeheer Regio Oost for allowing permission to work in Millingerwaard and Ciska Raaijmakers for invaluable help during the experiment.

LITERATURE CITED

- Agrawal AA, Kotanen PM, Mitchell CE, Power AG, Godsoe W, Klironomos J. 2005. Enemy release? An experiment with congeneric plant pairs and diverse above- and belowground enemies. *Ecology* 86:2979-2989.
- Andow DA, Imura O. 1994. Specialization of phytophagous arthropod communities on introduced plants. *Ecology* 75:296-300.
- Bever JD, Westover KM, Antonovics J. 1997. Incorporating the soil community into plant population dynamics: the utility of the feedback approach. *Journal of Ecology* 85:561-573.
- Bezemer TM, Fountain MT, Barea JM, Christensen S, Dekker SC, Duyts H, van Hal R, Harvey JA, Hedlund K, Maraun M, Mikola J, Mladenov AG, Robin C, de Ruiter PC, Scheu S, Setälä H, Smilauer P, van der Putten WH. 2010. Divergent composition but similar function of soil food webs of individual plants: plant species and community effects. *Ecology* 91: 3027-3036.
- Brinkman EP, Van der Putten WH, Bakker E-J, Verhoeven KJF. 2010. Plant-soil feedback: experimental approaches, statistical analyses and ecological interpretations. *Journal of Ecology* 98:1063-1073.

- Bucharova A, van Kleunen M. .2009. Introduction history and species characteristics partly explain naturalization success of North American woody species in Europe. *Journal of Ecology* 97:230-238
- Callaway RM, Thelen GC, Rodriguez A, Holben WE. 2004. Soil biota and exotic plant invasion. *Nature* 427:731-733.
- Carpenter D, Cappuccino N. 2005. Herbivory, time since introduction and the invasiveness of exotic plants. *Journal of Ecology* 93:315-321.
- Chun YJ, van Kleunen M, Dawson W. 2010. The role of enemy release, tolerance and resistance in plant invasions: linking damage to performance. *Ecology Letters* 13:937-946.
- Diez JM, Dickie I, Edwards G, Hulme PE, Sullivan JJ, Duncan RP. 2010. Negative soil feedbacks accumulate over time for non-native plant species. *Ecology Letters* 13:803-809
- Dirkse GM, Hochstenback SMH, Reijerse AI, Bijlsma R-J, Cerff D. 2007. *Flora van Nijmegen en Kleef 1800-2006: Catalogus van soorten met historische vindplaatsen en recente verspreiding*. Mook
- Dostál P, Müllerová J., Pyšek P., Pergl J. & Klinerová T. 2013. The impact of an invasive plant changes over time. *Ecology Letters* 16: 1277–1284.
- Elton C. 1958. *The ecology of invasions by enemies and plants*.
- Engelkes T, Morriën E, Verhoeven KJF, Bezemer TM, Biere A, Harvey JA, McIntyre LM, Tamis WLM, van der Putten WH. 2008. Successful range-expanding plants experience less above-ground and below-ground enemy impact. *Nature* 456:946-948
- Firn J, Moore JL, MacDougall AS, Borer ET, Seabloom EW, HilleRisLambers J, Harpole WS, Cleland EE, Brown CS, Knops JMH, Prober SM, Pyke

- DA, Farrell KA, Bakker JD, O'Halloran LR, Adler PB, Collins SL, D'Antonio CM, Crawley MJ, Wolkovich EM, La Pierre KJ, Melbourne BA, Hautier Y, Morgan JW, Leakey ADB, Kay A, McCulley R, Davies KF, Stevens CJ, Chu C-J, Holl KD, Klein JA, Fay PA, Hagenah N, Kirkman KP, Buckley YM. 2011. Abundance of introduced species at home predicts abundance away in herbaceous communities. *Ecology Letters* 14:274-281.
- Gange AC, Brown VK. 1989. Insect herbivory affects size variability in plant populations. *Oikos* 56:351-356.
- Gassó N, Sol D, Pino J, Dana ED, Lloret F, Sanz-Elorza M, Sobrino E, Vilà M. 2009. Exploring species attributes and site characteristics to assess plant invasions in Spain. *Diversity and Distributions* 15:50-58
- Genung MA, Bailey JK, Schweitzer JA. 2012. Welcome to the neighbourhood: interspecific genotype by genotype interactions in *Solidago* influence above- and belowground biomass and associated communities. *Ecology Letters* 15:65-73.
- Grayston SJ, Wang SQ, Campbell CD, Edwards AC. 1998. Selective influence of plant species on microbial diversity in the rhizosphere. *Soil Biology & Biochemistry* 30:369-378.
- Gundale MJ, Kardol P, Nilsson MC, Nilsson U, Lucas RW and Wardle D A. 2014. Interactions with soil biota shift from negative to positive when a tree species is moved outside its native range. *New Phytologist*, 202: 415–421.

- Hamilton MA, Murray BR, Cadotte MW, Hose GC, Baker AC, Harris CJ, Licari D. 2005. Life-history correlates of plant invasiveness at regional and continental scales. *Ecology Letters* 8:1066-1074
- Hawkes CV. 2007. Are invaders moving targets? The generality and persistence of advantages in size, reproduction, and enemy release in invasive plant species with time since introduction. *American Naturalist* 170:832-843.
- Johnson DJ, Beaulieu WT, Bever JD, Clay K. 2012. Conspecific Negative Density Dependence and Forest Diversity. *Science* 336:904-907.
- Keane RM, Crawley MJ. 2002. Exotic plant invasions and the enemy release hypothesis. *Trends in Ecology & Evolution* 17:164-170
- Klironomos JN. 2002. Feedback with soil biota contributes to plant rarity and invasiveness in communities. *Nature* 417:67-70
- Kowalchuk GA, Buma DS, de Boer W, Klinkhamer PGL, van Veen JA. 2002. Effects of above-ground plant species composition and diversity on the diversity of soil-borne microorganisms. *Antonie Van Leeuwenhoek International Journal of General and Molecular Microbiology* 81:509-520.
- Kulmatiski A, Beard KH, Stevens JR, Cobbold SM. 2008. Plant-soil feedbacks: a meta-analytical review. *Ecology Letters* 11:980-992.
- Lankau RA. 2011. Resistance and recovery of soil microbial communities in the face of *Alliaria petiolata* invasions. *New Phytologist* 189:536-548.
- Lankau RA, Nuzzo V, Spyreas G, Davis AS. 2009. Evolutionary limits ameliorate the negative impact of an invasive plant. *Proceedings of the*

National Academy of Sciences of the United States of America
106:15362-15367.

- Lau JA and Lennon JT. 2011. Evolutionary ecology of belowground-
aboveground interactions: soil microbial structure alters selection on
plant traits. *New Phytologist* 192:215-224.
- Lau JA and Lennon JT. 2012. Rapid responses of soil microorganisms
improve plant fitness in novel environments. *Proceedings of the
National Academy of Sciences*. 109:14058-14062.
- Levine JM, Pachepsky E, Kendall BE, Yelenik SG, Hille Ris Lambers J. 2006.
Plant-soil feedbacks and invasive spread. *Ecology Letters* 9:1005-
1014.
- Mangan SA, Schnitzer SA, Herre EA, Mack KML, Valencia MC, Sanchez EI,
Bever JD. 2010. Negative plant-soil feedback predicts tree-species
relative abundance in a tropical forest. *Nature* 466:752-U710.
- Maron JL, Vilà M. 2001. When do herbivores affect plant invasion? Evidence
for the natural enemies and biotic resistance hypotheses. *Oikos*
95:361-373
- Milbau A, Stout JC. 2008. Factors associated with alien plants transitioning
from casual, to naturalized, to invasive. *Conservation Biology* 22:308-
317
- Mitchell CE, Blumenthal D, Jarosik V, Puckett EE, Pyšek P. 2010. Controls on
pathogen species richness in plants' introduced and native ranges:
roles of residence time, range size and host traits. *Ecology Letters* 13
12:1525-1535.

- Mitchell CE, Power AG. 2003. Release of invasive plants from fungal and viral pathogens. *Nature* 421:625-627. Müller-Schärer H, Schaffner U, Steinger T. 2004. Evolution in invasive plants: implications for biological control. *Trends in Ecology and Evolution* 19 :417-422
- Pyšek P, Richardson DM, Williamson M. 2004. Predicting and explaining plant invasions through analysis of source area floras: some critical considerations. *Diversity and Distributions* 10:179-187
- Reinhart KO. 2012. The organization of plant communities: negative plant–soil feedbacks and semiarid grasslands. *Ecology* 93:2377-2385.
- Reinhart KO, Packer A, Van der Putten WH, Clay K. 2003. Plant-soil biota interactions and spatial distribution of black cherry in its native and invasive ranges. *Ecology Letters* 6:1046-1050.
- Reinhart KO, Tytgat T, Van der Putten WH, Clay K. 2010. Virulence of soil-borne pathogens and invasion by *Prunus serotina*. *New Phytologist* 186:484-495.
- Schaminée JHJ, Hennekens SM, Ozinga WA. 2007. Use of the ecological information system SynBioSys for the analysis of large datasets. *Journal of Vegetation Science* 18:463-470
- Schweitzer JA, Bailey JK, Fischer DG, LeRoy CJ, Lonsdorf EV, Whitham TG and Hart SC. 2008. Soil microorganism-plant interactions: Heritable relationships between plant genotype and associated microorganisms. *Ecology* 89:773-781.
- Simberloff D, Gibbons L. 2004. Now you see them, now you don't - population crashes of established introduced species. *Biological Invasions* 6:161-172.

- Speek TAA, Lotz LAP, Ozinga WA, Tamis WLM, Schaminée JHJ, van der Putten WH. 2011. Factors relating to regional and local success of exotic plant species in their new range. *Diversity and Distributions* 17:542-551.
- Tamis WLM, van der Meijden R, Runhaar J, Bekker RM, Ozinga WA, Odé B, Hoste I. 2004. Standaardlijst van de Nederlandse flora 2003. *Gorteria* 30 :101-195
- van Grunsven RHA, van der Putten WH, Bezemer TM, Berendse F, Veenendaal EM. 2010. Plant-soil interactions in the expansion and native range of a poleward shifting plant species. *Global Change Biology* 16:380-385.
- van Grunsven RHA, van der Putten WH, Bezemer TM, Tamis WLM, Berendse F, Veenendaal EM. 2007. Reduced plant-soil feedback of plant species expanding their range as compared to natives. *Journal of Ecology* 95:1050-1057.
- van der Putten WH, Vandijk C, Peters BAM. 1993. Plant-specific soil-borne diseases contribute to succession in foredune vegetation. *Nature* 362:53-56
- Van der Putten, WH, Van Dijk C and Troelstra SR. 1988. Biotic soil factors affecting the growth and development of *Ammophila arenaria*. *Oecologia* 76: 313-320.
- Wilson JRU, Richardson DM, Rouget M, Proches S, Amis MA, Henderson L, Thuiller W. 2007. Residence time and potential range: crucial considerations in modelling plant invasions. *Diversity and Distributions* 13 :11-22

FIGURE LEGENDS

Figure 1. Mean soil feedback effect on the biomass of shoots and roots in relation to the residence time or the local dominance of naturalized exotic plant species in the Netherlands. Each circle represents a different plant species.




Figure 1. Mean soil feedback effect on the biomass of shoots and roots in relation to the residence time or the local dominance of naturalized exotic plant species in the Netherlands. Each circle represents a different plant species.